AD-A277 376 ATION PAGE

Form Appresen DAME NO COMO 188

of Management and Budget Papers in 1997

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE

3. REPORT TYPE AND DATES COVERED

30 Sep 92 TO 29 Sep 93 FINAL

4. TITLE AND SUBTITLE

(FY91 AASERT) NEUROTRANSMITTERS AND PHOSPHOLIPID

METABOLISM

6. AUTHOR(S)

DR RICHARD J. WURTMAN

5 FUNDING NUMBERS

F49620-92-J-0479

61103D

3484

S4

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Massachusetts Institute of Technology Dept of Brain & Cognitive Sciences 77 Massachusetts Avenue (E25-604) Cambridge, MA 02139

PERFORMING ORGANIZATION KUPORT NUMBER

AFOSR-TR-

0050 94

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

AFOSR/NL 110 Duncan Avenue, Suite Bl15 Bolling AFB DC 20332-0001

10. SPONSGRING MONITORING AGENCY REPORT NUMBER

Dr Haddad

11. SUPPLEMENTARY NOTES

12a, DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited

12. AESTRACT (Maximum 200 words)

After receiving the AASERT fellowship, I conducted post-nBM lesion timecourse studies in order to determine whether cholinergic and phospholipid metabolism recovers after unilateral lesions. I found that cortical phospholipid biosynthesis and hydrolysis is altered following loss of cholinergic input.



7 1

. 14 SUBJECT TERMS 5 NUMBER OF PAGES 16 PRICE CODE SECURITY CLASSIFICATION 18. SECURITY CLASSIFICATION SECURITY CLASSIFI ATION LIME AT IN OF ABSTRACT OF ABSTRACT OF REPORT OF THIS PAGE (UL) (U) (U) (U)

1.11 F40 01 280 5500







Approved a militarelease; district

Progress Report for AFOSR/AASERT 2/16/94

Todd C. Holmes, Ph.D., Fellowship Recipient

The offices of lesion of ascending modulatory transmitter systems on chospholipid metabolism in the rat brain: models for neurotransmitter regulation of phospholipid metabolism.

My thesis research in Dr. Richard Wurtman's laboratory centered on the interactions between acetylcholine and phospholipid metabolism, specifically, on changes of phospholipid metabolism and mass in cortical terminal fields after removal of cholinergic innervation. Before receiving the AASERT fellowship, I found that phospholipid and phospholipid metabolite levels in frontoparietal cortex are decreased following excitotoxic lesion of the basal forebrain cholinergic nucleus basalis (nBM). nBM lesions were verified by the measurement of choline acetyltransferase activity and evoked acetylcholine release in cortical slices.

After receiving the AASERT fellowship, I conducted post-nBM lesion timecourse studies in order to determine whether cholinergic and phospholipid metabolism recovers after unilateral lesions. Choline acetyltransferase activity and evoked acetylcholine release in cortical slices showed an eventual recovery after the lesion. The decreases in cortical phospholipids also recovered after the lesion, the time course was similar to that seen for evoked acetylcholine release. Cholinergic lesion by surgical sectioning resulted in decreased cortical phospholipids, also excitotoxic lesions of non-cholinergic projections to cortex did not change phospholipid levels. Radiolabelling and enzyme assay experiments with cortical tissues indicate that phospholipid biosynthesis is decreased following nBM lesion. This decrease occurs at the point of choline kinase. Phospholipid turnover in contex may also be increased following nBM lesions, as indicated by muscarinit inositol lipid hydrolysis and choline radiolabelling of cortical tissues,

En conclusion, cortical phospholipid biosynthesis and hydrolysis is altered following loss of cholinergic input.

Dist Special

A-1

TOTAL P.82